

Quantum chromodynamics with various number of flavors *

Y. Iwasaki,^a K. Kanaya,^a S. Sakai^b and T. Yoshié^a

^aInstitute of Physics, University of Tsukuba, Ibaraki 305, Japan

^bFaculty of Education, Yamagata University, Yamagata 990, Japan

The phase structure of QCD with various number of flavors is studied for Wilson quarks. For the case of $N_F = 3$ we find that the finite temperature deconfining transition is of first order in the chiral limit on an $N_t = 4$ lattice. Together with our previous results that the deconfining transition in the chiral limit is continuous for $N_F = 2$ and is first order for $N_F = 6$, the order of the transition is found to be consistent with a prediction of universality. The case of $SU(2)$ QCD is also studied in the strong coupling limit and the phase structure is found to be quite similar to the case of $SU(3)$: There exists a critical number of flavors N_F^* and for $N_F \geq N_F^*$ the confinement is broken even in the strong coupling limit for light quarks. $N_F^* = 3$ corresponding to 7 for $SU(3)$.

1. INTRODUCTION

The deconfining transition of QCD is the last phase transition which our Universe has enjoyed in the past. To study the nature of this transition it is essential to include dynamical quarks. In this paper we study the phase structure of finite temperature lattice QCD with degenerate N_F Wilson quarks.

Our model has 4 parameters: N_F (the number of flavors), β (gauge coupling), K (hopping parameter) and N_t (lattice size in the temporal direction). For $N_t = 4 - 8$ we use a spatial $8^2 \times 10$ lattice (the lattice size 10 is doubled for hadrons), while for $N_t = 18$ we use $18^2 \times 24$. The method of simulation is similar to that described in [1,2].

1.1. $N_F = 2$ and 6

In a previous paper [1] we studied the deconfining transition K_T for the cases of $N_F = 2$ and 6 on an $N_t = 4$ lattice. Fig. 1 shows our result of the phase diagram for $N_F = 2$ near the chiral limit K_C . (The result for $N_F = 3$ will be discussed later.) K_T is identified by a sudden change in the behavior of physical observables such as plaquette and m_π^2 . K_C is determined by a linear extrapolation (in $1/K$) of m_π^2 in the confining region on the lattice with the same N_t . We find that N_t - (as well as N_F -) dependence of K_C is small for $\beta \lesssim 4.5$ compared with the difference of




Figure 1. Phase diagram for $N_F = 2$ and 3 on an $N_t = 4$ lattice.

K_C determined by m_π^2 and m_q ; the difference is a result of the $O(a)$ corrections concerning the chiral behavior of Wilson fermions. The location of the crossing point β_{CT} is identified by monitoring N_{inv} , the number of CG iterations to invert the quark matrix, just on the K_C -line [1,2]. This can be done because of an empirical law [3] that N_{inv} becomes very large in the confining phase when we get close to K_C , while N_{inv} remains small in the deconfining phase even at K_C .

The nature of the transition at β_{CT} can be studied by measuring observables on the K_C -line [1]. We find that, when we decrease β toward

*Talk presented by K. Kanaya at *Lattice 93*, Dallas, USA.




Figure 2. (a) m_π^2 and $2m_q$ and (b) N_{inv} history for $N_F = 3$ on K_C -line. Jackknife errors for (a) are smaller than the symbols.

β_{CT} , m_π^2 decreases linearly and is consistent with zero at β_{CT} . This implies that the transition is continuous (second order or crossover), in accord with the result of an universality argument by Rajogopal and Wilczek who predict the second order transition for $N_F = 2$ [4].

This result is in clear contrast with the case of $N_F = 6$ where we find that m_π^2 is large near and above $\beta_{CT} \simeq 0.3$. We also find a two-state signal in the time development of N_{inv} and other observables at $\beta = 0.3$ [1]. The first order transition observed is again consistent with a prediction of an universality argument stating that the transition is of first order for $N_F \geq 3$ [5].

2. $N_F = 3$

Because the effective N_F at the deconfining transition temperature in the real world is between 2 and 3, it is important to extend the study to the case of 3 flavors. Performing simulations

we find the phase diagram shown in Fig. 1. As expected, K_T shifts to smaller β with increasing N_F from 2 to 3. Fig. 2 (a) is the result for m_π^2 on the K_C line. We find that m_π^2 is large for $\beta \gtrsim \beta_{CT} \simeq 3.0$. We also find a two-state signal at $\beta = 3.0$ (Fig. 2 (b)). We therefore conclude that the deconfining transition is of first order for $N_F = 3$ in the chiral limit, which is in accord with the prediction of universality by Pisarski and Wilczek [5].

3. MANY FLAVORS

3.1. $SU(3)$ case

Now let us consider the case of much larger N_F . In our previous paper we have shown that, when $N_F \geq 7$, light quarks are not confining even in the strong coupling limit[2]. Extending the study to finite β we find the following phase structure for $N_F \geq 7$ [1]: We have a N_t -independent deconfining transition line K_D , which starts from a finite K at $\beta = 0$ and extends to larger β . In addition, we have a N_t -dependent deconfining transition/crossover line K_T , which reaches the K_D -line without crossing the K_C -line.

To understand this phase structure, let us recall the property of the perturbative β function. Asymptotic freedom is broken for $N_F \geq 17$ because the first coefficient of the β function becomes positive. We note that the second coefficient changes its sign for $N_F \geq 9$. This suggests the existence of an infrared fixed point (IRFP) for $16 \geq N_F \geq N_F^*$ ($N_F^* = 9$ up to 2-loop perturbation theory). With an IRFP correlation functions scale with non-canonical powers at large distances and therefore there are no hadronic particles[6]. Our MC result [1,2] implies that $N_F^* = 7$.

3.2. $SU(2)$ case

As an extension of the $SU(3)$ case, we study an $SU(2)$ QCD in the strong coupling limit. We note that the $SU(2)$ β -function has the same characteristics as $SU(3)$ except for that typical numbers for N_F are generally smaller than those for $SU(3)$: With $SU(2)$, asymptotic freedom is lost when $N_F \geq 11$ and 2-loop result for N_F^* is 6. This is natural because the confining force is weaker for $SU(2)$. We may therefore expect that N_F^* is




Figure 3. Observables in $SU(2)$ QCD for $K = 0.25$ at $\beta = 0$ as a function of N_F : (a) plaquette and Polyakov loop and (b) m_π^2 and $2m_q$.

smaller than the case of $SU(3)$.

Performing a simulation for $N_F = 8$ and $N_t = 4$ we find that the deconfining transition appears at $K = 0.2 - 0.21$ and that the chiral limit at $K = 0.25$ is in the deconfining phase. Decreasing N_F gradually with fixing $K = K_C$, we find that the chiral limit remains in the deconfining phase down to $N_F = 3$. When we further decrease N_F to 2, N_{inv} shows a rapid increase with the molecular-dynamics time and finally exceeds 10000 in clear contrast with small numbers of $O(10^2)$ for $N_F \geq 3$. The chiral limit is therefore in the confining phase for $N_F = 2$ [2]. Fig. 3 shows our results for physical observables for $K = K_C$. To study the N_t dependence of the transition, we simulate on an $N_t = 8$ lattice for the critical case of $N_F = 3$. The stability of the result confirms that the deconfining transition we observe is a bulk transition. According to our experience with $SU(3)$ QCD, we expect that these properties in

the strong coupling limit hold also for small β . We therefore conclude that $N_F^* = 3$ for $SU(2)$ QCD.

4. CONCLUSION

We studied the phase structure of QCD with dynamical Wilson quarks. Extending our previous study, we found that the finite temperature deconfining transition for $N_F = 3$ is of first order. We also studied an $SU(2)$ QCD and found that N_F -dependence of $SU(2)$ QCD is quite similar to that of $SU(3)$, but now N_F^* , above which confinement is lost for light quarks, is 3 instead of 7 for $SU(3)$.

We have studied the phase structure of QCD mainly on an $N_t = 4$ lattice. To study the implication to the continuum limit, the system should be investigated also at larger β on a larger lattice [7,8]. We are extending our study in this direction.

The simulations are performed with HITAC S820/80 at KEK and with QCDPAX. We thank members of KEK for their support and the other members of QCDPAX collaboration for their help. This project is in part supported by the Grant-in-Aid of Ministry of Education, Science and Culture (No.62060001 and No.02402003).

REFERENCES

1. Y. Iwasaki, K. Kanaya, S. Sakai, and T. Yoshié, Nucl. Phys. B (Proc. Suppl.) 30 (1993) 327; *ibid.* 26 (1992) 311.
2. Y. Iwasaki, K. Kanaya, S. Sakai, and T. Yoshié, Phys. Rev. Lett. 69 (1992) 21.
3. Y. Iwasaki, K. Kanaya, S. Sakai, and T. Yoshié, Phys. Rev. Lett. 67 (1991) 1494.
4. K. Rajagopal and F. Wilczek, Nucl. Phys. B399 (1993) 395.
5. R. Pisarski and F. Wilczek, Phys. Rev. D29 (1984) 338.
6. T. Banks and A. Zaks, Nucl. Phys. B196 (1982) 189.
7. C. Bernard, *et al.*, Phys. Rev. D46 (1992) 4741; preprint AZPH-TH/93-29; D. Tous-saint, this proceedings.
8. D. Zhu and N. Christ, this proceedings.